

Category	Brief Description	Text
ATRIAL FIBRILLATION	Monthly INR	<b>Monthly INR</b> Number of calendar months in which at least one INR measurement was made.
ATRIAL FIBRILLATION	Chronic anticoagulation therapy	<b>Chronic anticoagulation therapy</b> Patients who were prescribed warfarin (or other anticoagulant) during the 12-month reporting period.
CORONARY ARTERY DISEASE	Symptom control	<b>Symptom control</b> Percentage of visits for patients aged 50 years and older with a diagnosis of CAD who are angina-free OR are prescribed at least two anti-anginal medications.
CORONARY ARTERY DISEASE	Symptom and activity assessment	<b>Symptom and activity assessment</b> Percentage of patients aged 50 years and older with a diagnosis of CAD who were evaluated for both level of activity and anginal symptoms during one or more visits.
CORONARY ARTERY DISEASE	Oral antiplatelet therapy for patients with CAD	<b>Oral antiplatelet therapy for patients with CAD</b> Percentage of patients aged 50 years and older with a diagnosis of CAD who were prescribed aspirin or clopidogrel
CORONARY ARTERY DISEASE	Blood pressure treatment	<b>Blood pressure treatment</b> Percent of patients less than 60 years old AND BP greater than or equal to 140/90 AND who are taking or were prescribed two or more antihypertensive agents at most recent visit during the previous 12 months.
CORONARY ARTERY DISEASE	Blood pressure control	<b>Blood pressure control</b> Percent of patients < 60 with blood pressure < 140/90.
CORONARY ARTERY DISEASE	Beta-blocker therapy for CAD patients with prior MI	<b>Beta-blocker therapy for CAD patients with prior MI</b> Percentage of patients aged 18 years and older with a diagnosis of CAD and prior MI who were prescribed beta-blocker therapy.
CORONARY ARTERY DISEASE	ACE or ARB therapy for patients with CAD AND (diabetes and/or LVSD)	<b>ACE or ARB therapy for patients with CAD, diabetes and/or LVSD</b> Percentage of patients who also have diabetes mellitus and/or LVSD (LVEF < 40%) who were prescribed ACE inhibitor or ARB therapy. (PQRS-comparable)
DIABETES	Urine screening for microalbumin or medical attention for nephropathy	<b>Urine screening for microalbumin or medical attention for nephropathy</b> Percentage of patients aged 50 through 75 years with diabetes who received urine protein screening or medical attention for nephropathy during at least one office visit within 12 months. (PQRS-comparable)

Category	Brief Description	Text
DIABETES	High blood pressure control (DM)	<b>High blood pressure control (DM)</b> Percentage of patients aged 50 through 75 years with diabetes who had most recent blood pressure in control (less than 140/90 mmHg).
DIABETES	Hemoglobin A1c control	<b>Hemoglobin A1c control</b> Percentage of patients aged 50 through 75 years with diabetes who had the following values for their most recent hemoglobin A1c: < 7.0
DIABETES	Foot exam	<b>Foot exam</b> The percentage of patients aged 18 through 75 years with diabetes who had a foot examination. (PQRS-comparable)
DIABETES	Dilated eye exam	<b>Dilated eye exam</b> Percentage of patients aged 50 through 75 years with diabetes who had a dilated eye exam.
HEART FAILURE	Left ventricular function (LVF) assessment	<b>Left ventricular function (LVF) assessment</b> Percentage of patients aged 50 years and older with a diagnosis of heart failure who have quantitative or qualitative results of LVF assessment recorded.
HYPERTENSION	Blood pressure control (hypertension)	<b>Blood pressure control</b> Percentage of patients ages 50-59 with BP< 140/90
HYPERTENSION	Blood pressure control (hypertension)	<b>Blood pressure control</b> Percentage of patients ages > = 60 with BP< 150/90
PREVENTIVE CARE AND SCREENING	Unhealthy alcohol use - screening	<b>Unhealthy alcohol use - screening</b> Percentage of patients aged 50 years and older who were screened for unhealthy alcohol use using a systematic screening method within 24 months.
PREVENTIVE CARE AND SCREENING	Pneumonia vaccination for smokers	<b>Pneumonia vaccination for smokers</b> Percentage of active smokers who have ever received a pneumococcal vaccine.
PREVENTIVE CARE AND SCREENING	Inquiry regarding tobacco use	<b>Inquiry regarding tobacco use</b> Percentage of patients who have been screened for tobacco use.
PREVENTIVE CARE AND SCREENING	Influenza immunization for patients > = 50 years old	<b>Influenza immunization for patients &gt; = 50 years old</b> Percentage of patients aged 50 years and older who received an influenza immunization during the flu season (September through February).
PREVENTIVE CARE AND SCREENING	Body Mass Index (BMI) - screening and follow-up	<b>Body Mass Index (BMI) – screening and follow-up</b> Percentage of patients with a documented calculated BMI in the past six months outside of parameters who have a documented follow-up plan

Category	Brief Description	Text
PREVENTIVE CARE AND SCREENING	BMI control	<b>BMI control</b> Percent of patients with a calculated BMI in last 6 months outside of parameters
PREVENTIVE CARE AND SCREENING	Blood lipid therapy and control	<b>Blood lipid therapy and control</b> Proportion of patients who are taking appropriate statin therapy.
PREVENTIVE CARE AND SCREENING	Aspirin use in patients without clinical evidence of atherosclerotic disease who are at higher CVD risk	<b>Aspirin use in patients without clinical evidence of atherosclerotic disease who are at higher CVD risk</b> Patients who were advised to use aspirin.
PREVENTIVE CARE AND SCREENING	Advising smokers to quit	<b>Advising smokers to quit</b> Percentage of smokers who received a smoking cessation intervention in the last 24 months